



La CLAVE (The Clue or Key) A Psychoeducational Tool to Reduce Treatment Delay in Latinx with First Episode Psychosis



Many Latinos with serious mental illness (SMI) and their family members fail to recognize the signs of a first episode of psychosis symptoms. In general, they have poor psychosis literacy which may lead to longer delays in treatment and longer Duration of Untreated Psychosis (DUP).¹ López and colleagues developed a psychoeducational tool known as *La CLAVE* to reduce DUP among Latinos.^{1,2} *La CLAVE* was informed by conceptual models of health literacy.³

La CLAVE is a tool in Spanish that represents the symptoms of psychosis.²

C - false beliefs or delusions (*Creencias falsas*)


L - disordered speech or thought disorder (*Lenguaje desorganizado*)

A - hallucinations (*Alucinaciones*)

v - seeing things that others do not see (*Ver cosas que otros no ven*)

e - hearing sounds or voices that others do not hear (*Escuchar voces que no existen*)

La CLAVE uses plain language to guide individuals and caregivers to recognize the three domains of psychosis literacy (i.e., knowledge of psychosis, attributions to mental illness, and help-seeking behaviors).¹ *La CLAVE* uses popular Latin American music, video clips, or art. The purpose of using this guide with visual arts is to increase participant's knowledge and understanding of psychosis and self-efficacy skills to recognize the symptoms of psychosis in other individuals.^{1,2,4,5}



Conceptual Model of Psychosis Literacy (image)

By identifying these symptoms, individuals and their caregivers will be able to detect SMI earlier and seek treatment. La CLaVe reduces the duration of untreated psychosis (DUP).⁶ DUP is associated with worse long-term functioning because they can present serious symptoms, decreased remission rates, poorer social functioning, and less treatment benefits.^{7,8} DUP has been attributed to several reasons, including not recognizing SMI symptoms due to low mental health literacy.⁹

For more information visit: <https://www.uselaclave.com> and <https://youtu.be/Nb0jsnp4bP8>



@hilaMHTTC

hispaniclatino@mhttcnetwork.org

<https://mhttcnetwork.org/hispaniclatino>

For information contact us

(787) 785-5220

References

1. Lopez, S. R., Gamez, D., Mejia, Y., Calderon, V., Lopez, D., Ullman, J., & Kopelowicz, A. (2018). Psychosis literacy of U.S. Latinos with first episode psychosis and their caregivers. *Psychiatric Services*, 69(11), 1153-1159. doi: 10.1176/appi.ps.201700400
2. Lopez, S. R., Lara, M.C., Kopelowicz, A., Solano, S., Lara, M., Focerrada, H., & Aguilera, A. (2009). La CLaVe to increase psychosis literacy of Spanish-speaking community residents and family caregivers. *Journal of Consulting and Clinical Psychology*, 77(4), 763- 774. doi: 10.1037/a0016031
3. Institute of Medicine. (2004). *Health literacy: A prescription to end confusion*. National Academies Press. doi: 10.17226/10883
4. Casas, R. N., Gonzales, E., Aldana-Aragon, E., Lara-Muñoz, M., Kopelowicz, A., Andrews, L., & Lopez, S. (2014). Toward the early recognition of psychosis among Spanish speaking adults on both sides of the US-Mexico border. *Psychological Services*, 11(4), 460-469. <https://psycnet.apa.org/doi/10.1037/a0038017>
5. Calderon, V., Mejia, Y., Lara-Muñoz, M. del C., Segoviano, J., Castro, Q., & Lopez, S.R. (2015). Towards the sustainability of information campaigns: Training promotores to increase the psychosis literacy of Spanish-speaking communities. *Social Psychiatry and Psychiatric Epidemiology*, 50(4), 665-669. doi: 10.1007/s00127-014-0992-z
6. Melle, I., Larsen, T. K., Haahr, U., Friis, S., Johannessen, J. O., Opjordsmoen, S., ... & McGlashan, T. (2004). Reducing the duration of untreated first- episode psychosis: Effects on clinical presentation. *Archives of General Psychiatry*, 61(2), 143-150. doi: 10.1001/archpsyc.61.2.143
7. Penttila, M., Jasskelainen, E., Hirvonen, N., Isohanni, M., & Miettunen, J. (2014). Duration of untreated psychosis as predictor of long-term outcome in schizophrenia: systematic review and meta-analysis. *British Journal of Psychiatry*, 205(2), 88-94. doi: 10.1192/bjp.bp.113.127753.
8. Kane, J.M., Schooler, N.R., Marcy, P., Correll, C.U., Brunette, M.F., Mueser, K.T. ... & Robinson, D.G. (2015). The RAISE early treatment program for first-episode psychosis: Background, rationale, and study design. *Journal of Clinical Psychiatry*, 76(3), 240–246. doi: 10.4088/JCP.14m09289
9. Marshall, M., Lewis, S., Lockwood, A., Drake, R., Jones, P., & Croudace, T. (2005). Association between duration of untreated psychosis and outcome in cohorts of first-episode patients: a systematic review. *Archive of General Psychiatry*, 62(9), 975–83. doi: 1001/ archpsyc.62.9.975.